

and much lamented Commodore Goodenough, who met his death whilst out with Messrs. Perry and Corrie on an entomological excursion.

Family PAPILIONIDÆ.

Genus BELENOIS, Hübner.

10. BELENOIS NISEIA, McLeay (one male).

Havannah Harbour, Waté or Sandwich Island.

The single example sent differs a little from the Australian type, inasmuch as it resembles *B. clytie* ♂ on the upperside; but, without seeing more specimens, it would be unsafe to consider it a distinct species.

7. Descriptions of new Birds obtained by Mr. C. Buckley in Bolivia. By P. L. SCLATER, M.A., Ph.D., F.R.S., and OSBERT SALVIN, M.A., F.R.S.

[Received January 14, 1876.]

A collection of bird-skins formed by Mr. C. Buckley during his recent expedition into Bolivia, at several localities on the eastern slope of the main chain of the Andes, contains about 250 specimens, referable to 133 species. As a second collection is shortly expected from Mr. Buckley, we reserve a full account of the first one until the series is complete, and for the present only give descriptions of the new species which it contains. These are:—

1. DIGLOSSA GLAUCA, sp. nov.

Ex cinereo cærulescens fere unicolor; alis caudaque nigris cæruleo limbatis: subtus obscurior, remigum marginibus internis cinereis: rostro corneo, mandibula inferiore ad basin alba, pedibus nigris: long. tota 4·0, alæ 2·3, caudæ 1·6.

Hab. Nairapi, Bolivia (*Buckley*).

Mus. S.-G.

Of this new *Diglossa* Mr. Buckley sends but one specimen. It is a small species belonging to the "homochroous" section of Sclater's arrangement (*Ibis*, 1875, p. 207), and may be best placed between *D. plumbea* and *D. personata*. From the latter it is distinguishable by its much inferior size, from the former by its colour.

2. BUARREMON MELANOPS, sp. nov.

Buarremon rufinucha, Sclater, Cat. A. B. p. 91 (nec Lafr. et D'Orb.).

Supra niger; pileo cum nucha castaneo; lateribus capitis et mento summo nigris: subtus flavus, hypochondriis et crisso in olivaceum transeuntibus: rostro nigro, pedibus obscure corneis: long. tota 6·0, alæ 2·7, caudæ 2·6.

Hab. Simacu, Bolivia (*Buckley*).

Mus. S.-G. et P. L. S.

Obs. B. rufinucha (Lafr. et D'Orb.) * macula frontali utrinque sulphurea sane species diversa est.

Slater has long had two examples of this species in his collection obtained from the Maison Verreaux. They were referred to *B. rufinucha* erroneously, as it appears on examination of the specimens of that species in the British Museum. Mr. Buckley sends us five examples of this bird, all obtained at Simacu.

3. LEPTOPOGON TRISTIS, sp. nov.

Supra olivaceus, unicolor; alis nigris, tectricum alarium et secundariorum externorum apicibus macula quadrata sulphureo-alba ornatis; remigum et rectricum marginibus externis olivaceis: subtus sulphureo-flavus: subalaribus flavis: rostro et pedibus nigris: long. tota 4.4, alæ 2.15, caudæ 2.0, tarsi 0.75.

Hab. Bolivia, Simacu (*Buckley*).

Mus. S.-G. et P. L. S.

Obs. Sp. corporis colore supra olivaceo, subtus omnino flavo, ab aliis hujusce generis satis definita.

Slater has long had a single worn and imperfect skin of this *Leptopogon* in his collection under the MS. name which we now publish. Mr. Buckley sends but a single specimen. Slater's example was obtained from a dealer in Paris, without any locality attached to it.

4. HYPOXANTHUS ATRICEPS, sp. nov.

Hypoxanthus rivolii, Sel. et Salv. P. Z. S. 1873, p. 780.

Hab. Peruvia alta, Husampilla (*Whitely*); Bolivia, Ramosani (*Buckley*).

Mus. S.-G.

Obs. Similis *H. rivolii*, sed pileo summo in mari nigerrimo nec rubro.

An examination of a series of skins of this Woodpecker from Columbia, Ecuador, Peru, and Bolivia seems to show the form from the two latter countries is distinct—the head in the male being black, and never acquiring the brilliant crimson of the northern form.

February 15, 1876.

Prof. Mivart, F.R.S., in the Chair.

The following report on the additions to the Society's Menagerie during the month of January 1876 was read by the Secretary:—

The total number of registered additions to the Society's Menagerie during the month of January 1876 was 52, of which 32 were by presentation, 10 by purchase, and 10 were received on deposit. The total number of departures during the same period by death and removals was 78.

* Vide Slater, Syn. Av. Tanagr. p. 25.